

2018-08

Short-term interactive effects of increased temperatures and acidification on the calcifying macroalgae *Lithothamnion crispatum* and *Sonderophycus capensis*

Munoz, PT

<http://hdl.handle.net/10026.1/12304>

10.1016/j.aquabot.2018.04.008

Aquatic Botany

Elsevier

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

This is the author's accepted manuscript. The final published version of this work (the version of record) is published by Elsevier in Aquatic Botany made available online on the 26 April 2018 at: <https://doi.org/10.1016/j.aquabot.2018.04.008>. This work was made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

Short-term interactive effects of increased temperatures and acidification on the calcifying macroalgae *Lithothamnion crispatum* and *Sonderophycus capensis*

Pamela T. Muñoz^{1,2,3}, Claudio A. Sáez^{1*}, María Brezo Martínez-Callejas⁴, María Rosa Flores-Molina¹, Eduardo Bastos³, Alessandra Fonseca³, C. Frederico D. Gurgel³, José Bonomi Barufi³, Leonardo Röhrig³, Jason M. Hall-Spencer⁵, Paulo Antunes Horta³

1 Laboratory of Coastal Environmental Research, Centro de Estudios Avanzados, Universidad de Playa Ancha, Viña del Mar, Chile.

2 Programa de Doctorado Interdisciplinario en Ciencias Ambientales, Facultad de Ciencias Naturales y Exactas, Universidad de Playa Ancha, Valparaíso, Chile.

3 Programa de Pós-graduação em Biologia de Algas, Plantas e Fungos, Departamento de Botânica, Universidade Federal de Santa Catarina, Trindade, 88010–970, Florianópolis, Santa Catarina, Brazil.

4 Unidad de Biodiversidad y Consevación, Universidad Rey Juan Carlos, 28933 Mostoles, Spain.

5 Marine Biology and Ecology Research Centre, Plymouth University, Drake Circus, Plymouth, PL4 8AA United Kingdom

*Corresponding author: claudio.saez@upla.cl

Abstract

Combined effect of acidification and sea warming derived from future conditions of Climate Change have been little investigated in marine photoautotrophs, especially on sensitive organisms such as calcifying macroalgae. The aim of this investigation was to evaluate the interactive effects of acidification and increased temperatures on the two Brazilian calcifying macroalgae *Lithothamnion crispatum* and *Sonderophycus capensis*. Both species were cultured for 48 h under: (1) current pH (7.8 ± 0.2) and temperature ($18 \pm 2^\circ\text{C}$) during winter; (2) future pH (7.4 ± 0.2) and temperature ($30 \pm 2^\circ\text{C}$) during summer; (3) current temperature and future pH; and (4) future temperature and current

pH. We evaluated photosynthetic performance (measured $\Delta F/F'_m$), growth rates (weight), relative carbonate content, and total phenolic compounds. Our results showed similar negative effects under decreased pH and increased temperatures in both species, although carbonate content *S. capensis* was less affected than in *L. crispatum*. Total phenolic compounds measured in *S. capensis* showed the highest levels at potential future pH and temperature conditions. Given that stress conditions associated with decreased pH and increased temperatures are important inductors of an oxidative response, it is likely that phenolic compounds are synthesized to fulfil an antioxidant purpose. Even though physiological performance was affected in both calcifying macroalgae under the most likely negative future pH and temperature conditions, their biological viability indicates they may be able to thrive under coming Climate Change scenarios.

Keywords: Climate Change; acidification; warming; macroalgae.

INTRODUCTION

Atmospheric carbon dioxide (CO₂) concentrations have been increasing from 280 ppm since the pre-industrial period to above 400 ppm nowadays; moreover, CO₂ levels are expected to increase in up to 970 ppm by the end of the 21st Century (Harley et al. 2006; Harley et al. 2012). Predicted CO₂ rise and accompanied increase in global temperatures is expected to induce greater sea surface temperatures, which could fluctuate between 2 and 6°C depending on the strictness of the extrapolation (IPCC 2014). On the other hand, approximately 25% of atmospheric CO₂ is absorbed by the ocean, which in reaction with seawater increases the levels of carbonic acid (H₂CO₃), subsequently diminishing the pH; this process is expected to induce a 0.2-0.4 pH decline by the year 2100 (Feely et al. 2004). Ocean acidification produces changes in total dissolved inorganic carbon (DIC), manifested in higher concentration of carbonate ions (CO₃²⁻) and bicarbonate (HCO₃⁻); important molecules that may affect physiological processes, mostly in marine calcifying organisms (Martin and Hall-Spencer 2017; Orr et al. 2005; Koch et al. 2013). As signal of Climate Change, monitoring of coastal marine habitats has revealed a concerning increase in the incidence of mass mortalities in flora and fauna due to marine heat waves as observed in Australia (Wernberg et al. 2011) or in the south Atlantic (Ferreira et al.

2015). Elevated temperature, even over a short period, can be lethal due to biochemical damages (Gouvêa et al. 2017). In these coastal and shallow regions, high pH variability is observed, eventually exposing organisms to conditions frequently predicted to the end of this century considering ocean acidification (Duarte et al. 2013). Changes in ocean circulation also provide localized and short variability in ocean pH due upwelling/downwelling zones (Gruber et al. 2012; Findlay et al. 2013), CO₂ enrichment from volcanic vents (Hall-Spencer et al. 2008), and as a function of the metabolism of some marine communities (Anthony et al. 2011) or even in anoxic zones (Melzner et al. 2013). Therefore, extreme values of temperature and pH/pCO₂ present in different regions of the world can accentuate or even represent nowadays ocean acidification impacts on the biology of marine biodiversity.

Calcifying macroalgae are key ecological components of the Atlantic coast. They are considered ecosystem bioengineers, as create habitat, niche for settlement and nursery for other algae, invertebrates, and fish (Figueiredo et al. 2012; Riul et al. 2009; Amado-Filho et al. 2007). Furthermore, these organisms are major producers of carbonate sediment in the marine environment (Basso 2012). Previous studies have shown that unbalanced DIC mediated by decreased pH reduces the saturation of calcium carbonate (CaCO₃) in calcifying marine organisms, such calcifying macroalgae, corals, foraminifera and coccolithophores; this process can affect calcification rates when producing new skeleton (Basso 2012; Martin and Hall-Spencer 2017; Orr et al. 2005). CaCO₃ occurs in skeletons of calcifying macroalgae as calcite or aragonite; however, magnesium (Mg²⁺) can replace calcium in calcite macroalgae, which has been observed to be more prone to dissolution under acidification than macroalgae with aragonite skeleton (Basso 2012; Jury et al. 2010; Smith et al. 2012).

Calcifying macroalgae are capable of performing calcification and photosynthesis at different rates depending on CO₂ concentrations (Hofmann et al. 2012; Hurd et al. 2009; Koch et al. 2013). Despite that acidification can produce adverse effects on calcification rates, changing carbonate content, high concentrations of CO₂ have been detected to favor photosynthesis in some cases (Hurd et al. 2009; Koch et al. 2013). For instance, Semesi et al. (2009) showed that a progressive decline in pH induced a 13% increase in photosynthesis rates in coralline algae, although compromising their structure

due to 20% less calcification. Tropical and subtropical organisms survive near suboptimal physiological temperature; however, Climate Change-mediated increase in sea temperatures threatens their thermal tolerance thresholds (Latham 2008; Eggert 2012; Harley et al. 2006). Within tolerance ranges, rising temperature can induce an increase in photosynthesis, growth and carbonate deposition in calcifying macroalgae, but when these exceed tolerance thresholds, it can cause bleaching and chlorophyll degradation (Martin and Hall-Spencer 2017; Glynn 1996).

Drastic environmental disturbances can generate biological stress in macroalgae through the over-production of reactive oxygen species (ROS) (Bischof and Rautenberger 2012). ROS excess can produce damage by oxidizing lipids, proteins and nucleic acids; however, the cell can maintain ROS within homeostatic levels through the production of antioxidant compounds (Moenne et al. 2016). In this context, phenolic compounds have been observed to have strong ROS scavenging properties (Sáez et al. 2015; Flores-Molina et al. 2016). ROS-induced increase in phenolic compounds in macroalgae have been observed subject to several environmental stressors, among which can be mentioned herbivory, changes in temperature, salinity, irradiance and metal excess (Bischof and Rautenberger 2012; Moenne et al. 2016; Sáez et al. 2015; Flores-Molina et al. 2016).

Records on the interaction of different stressors associated with Climate Change on marine calcifying macroalgae are scarce; thus, more information on aspect such as the combined effects of acidification and increased temperatures would significantly improve our understanding on their current and future responses to progressive Climate Change. The aim of this research was evaluate the interactive effects of a short acidification and warming on physiological aspects of the calcifying red coralline *Lithothamnion crispatum* Hauck and the red aragonite *Sonderophycus capensis* (M) M.J. Wynne. *L. L. crispatum* has a wide range of latitudinal distribution; considered as a dominant species in the entire Atlantic coast of Brazil, its temperature tolerance ranges between 15 to 30°C (Pascelli et al. 2013; Riul et al. 2009). On the other hand, *S. capensis* has a limited distribution (southeastern Atlantic of Brazil) with tolerance ranges defined between 17 and 20°C (Zavialov et al. 1999). Taking into account limits of temperature tolerance and CaCO₃ types of skeleton in these species, it is expected that *L. crispatum* and *S. capensis*

have differential responses to single and combined effects of increased temperatures and acidification.

MATERIALS AND METHODS

Collection and experimental design. Entire individuals of two calcifying red macroalgae species, the coralline *L. crispatum* and the aragonite-based *S. capensis*, were collected at 10 m depth from Reserva Biologica do Arvoredo, in Santa Catarina (27° 15' S, 48° 20' W) and in Xavier Island (27° 36' S, 48° 23' O), respectively, in Brazil. After collection, individuals were stored in plastic bags with seawater, and immediately transported inside coolers to the Laboratory of Phycology at Universidad de Santa Catarina. Prior to experiments, the specimens were acclimated for 24 h with filtered seawater (0.45 µm) under constant aeration, temperature ($15 \pm 2^\circ\text{C}$) and light (50 µmol photons m⁻²s⁻¹). The experiment was carried out in a microcosm system over a short-term period (48 h). The experimental system consisted in two tanks with 50 L seawater that delivered seawater with a pump to 12 chambers of 1 L. Both macroalgae species were randomly distributed and maintained in four conditions: **(1)** average current pH (7.8 ± 0.2) and temperature ($18 \pm 2^\circ\text{C}$) registered in Florianopolis bay, Brazil, during winter **(2)**; future values of pH (7.4 ± 0.2) and temperature ($30 \pm 2^\circ\text{C}$) that can be extrapolated for Florianopolis bay during summer by the end of the 21st Century according to predictions by IPCC (2014); **(3)** Current temperature ($18 \pm 2^\circ\text{C}$) and future pH (7.4 ± 0.2); and **(4)** Future temperature ($30 \pm 2^\circ\text{C}$) and current pH (7.8 ± 0.2). For each treatment, three independent replicates were used. pH levels considered for control treatments respond to local levels registered in seawater nearby macroalgae collection sites. Certainly, in these locations there are several estuaries, such as Barra de Lagoa, that supply freshwater and inland material that induces a pH drop in coastal waters; in this context, several reports in this area show average pH levels around 7.8 (Martins-Pereira 2004; Dalinghaus 2016; Cancellier-Cechinel 2013). Considering the latter, acidification

treatments with a 0.4 pH decrease were made according IPCC predictions upon current local pH levels.

To achieve low pH conditions, bubbles of CO₂ were injected with a pump the experimental seawater. The added CO₂ were managed by a pH controller, model pH2010 (Weipro, China), connected to solenoid valves that regulated the amount of gas pumped into the seawater (Russell et al. 2009). Experimental temperatures were achieved by placing tanks on a homemade temperature-gradient table; this is a horizontal metal-made table, which surface temperature gradually increases from 15 °C to 35 °C, from one extreme to the other. In addition, the table constantly agitates to maintain temperature homogeneity in the seawater.

Water carbonate chemistry parameters. HCO₃⁻² and CO₃⁻² concentrations, *p*CO₂, calcite and aragonite saturation (Ω), were obtained via CO₂Sys_v2.1.xls program, using K1, K2, as in Millero et al. (1998). These values are summarized in Table 1. Estimations were based in measurements of pH, temperature, salinity and total alkalinity in seawater. Gran titration method was used to measure total alkalinity, using 0.0025 N hydrochloric acid (HCl; Merck) (Carmouze 1994); these measurements were taken after 48 h of experiments.

Photosynthetic performance. Photosynthetic measurements were performed throughout the course of the experiments in all treatments, at 6, 24 and 48 h. Effective quantum yield of photosystem II (PII; $\Delta F/F'_m$) was measured using a portable pulse-amplitude modulated (PAM) chlorophyll α fluorometer (DIVING PAM; Walz GmbH, Effeltrich, Germany). This parameter was measured according to Schreiber and Neubauer (Schreiber and Neubauer 1990): $\Delta F/F'_m = (F'_m - F_t)/F'_m$. In the latter, F'_m is the maximal fluorescence of light-acclimated sample induced by a saturating actinic light pulse (9000 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, 800 ms), and F_t is the intrinsic steady-state fluorescence emission in light-acclimated sample (Baker 2008).

Growth rate. To estimate the growth, the change in fresh weight (FW) biomass of the species was measured, prior and after 48 h of experiments. The mean relative growth was expressed as FW per day (FW d⁻¹), and calculated according to Packard and

192 Boardman (1999): $RG = (W_t - W_0)/t$. W_0 is the initial weight, W_t is final weight, and t is time
 193 in days (d).

194 **Relative carbonate content (RCC).** Biomass was dried at 60°C for 24 h and
 195 weighted. Initial weight (DW_i) was obtained and then immersed in 10% HCl (Merck) for
 196 24 h to initiate CaCO₃ dissolution, in order to maintain the same weight/volume ratio.
 197 After the experiments, samples were dried (60°C) again and were re-weighted (DW_f)
 198 (Figueroa et al. 2014). RCC as a proxy of calcification was calculated to the formula: %
 199 RCC ((DW_i-DW_f)/DW_f)*100).

200 **Content of total phenolic compound.** Due to biomass availability, total phenolic
 201 compounds were only determined in *S. capensis*, according to Randir et al. (2002). After
 202 48 h experiments, samples were immediately frozen in liquid nitrogen and stored at -
 203 80°C. Between 40-50 mg biomass were grounded to powder in a mortar with liquid
 204 nitrogen, at 4°C chamber in the darkness. One mL of 80% methanol (Merck) and 5 ml of
 205 95% methanol (Merck) were added to each sample in centrifuge tubes. Samples were
 206 centrifuged at 10,000 rpm for 5 min at room temperature. One mL of the supernatant was
 207 placed in a clean tube and mixed with 5 mL of distilled water, 1 mL of 95% ethanol and
 208 500 µL of Folin-Ciocalteu; the mixture was incubated for 5 min in the darkness. Then, 1
 209 mL of NaCO₃ (5% w/v) was added and incubated in the dark for 1 h at room temperature.
 210 Finally, absorbance was measured in a spectrophotometer, model P-220 (Biospectro,
 211 Brazil), at 725 nm. To calculate the content of total phenolic compounds from crude
 212 extracts, a standard curve of known concentrations of gallic acid was used (50 to 800 µg
 213 mL⁻¹ – $r^2 = 0.99$; $y = 1.254$).

214 **Statistical analyses.** Effects of temperature and pH on both species ($\Delta F/F_m'$,
 215 RCC, growth rate and phenolics content), and chemical changes in seawater, were
 216 assessed statistically by using a factorial analysis of variance (Factorial-ANOVA). A *post*
 217 *hoc* analysis of Student Newman-Keuls tests (SNK) was applied when Cochran's C test
 218 and visual inspection of residuals were conducted to test homogeneity of variances
 219 (Underwood 1997). A level of significance of 99% confidence interval ($p = 0.01$) was
 220 applied. These analyses were performed using the software Statistica version 7 (StatSoft
 221 Inc., Tulsa, OK, USA).

222 RESULTS

223 Photosynthetic responses

224 For both species, a significant interaction was found between temperature (18 and 30°C)
 225 and pH (7.8 and 7.4) on photosynthetic performance (measured as $\Delta F/Fm'$) (Factorial
 226 ANOVA, $p < 0.01$; Figure 1; Table 2). $\Delta F/Fm'$ significantly increased with respect to
 227 basal values in *L. crispatum* after 6, 24 and 48 h experiments at 18°C under both 7.4 and
 228 7.8 pH conditions (Figure 1A). A similar trend was observed in *L. crispatum* at 30°C
 229 under pH 7.4, although without significant differences at 24 h experiments (Figure 1A). Δ
 230 F/Fm' significantly decreased under pH 7.8 at 30°C in all experimental times, with lower
 231 values at 24 and 48 h experiments. In relation to *S. capensis*, no specific trends compared
 232 to basal values were observed at 18°C under both pH conditions and at 30°C under pH
 233 7.4 (Figure 1B). However, significantly lower $\Delta F/Fm'$ levels were recorded at 30°C
 234 under pH 7.8, especially at 24 and 48 h experiments (Figure 1; Table 2).

235 Responses in calcification

236 For both species a significant interaction was observed between temperature (18 and
 237 30°C) and pH (7.8 and 7.4) on RCC (Factorial ANOVA, $p < 0.01$; Figure 2; Table 2).
 238 Similar trends were observed in RCC between studied species. When decreasing the pH
 239 from 7.8 to 7.4, RCC decreased significantly under both 18 and 30°C (Figure 2). The
 240 exception was at 30°C in *L. crispatum*, where no significant differences were detected
 241 between both experimental pH conditions (Figure 2). }

242 Relative growth rate

243 No changes were recorded in growth rates in *L. crispatum*, with exception of the
 244 treatment at 30°C and pH 7.8, where negative values were observed (Figure 3A). A
 245 similar trend of negative growth rates were detected in *S. capensis* at 30°C and pH 7.8,
 246 although the rest of the treatments were similarly positive (Factorial ANOVA, $p < 0.01$;
 247 Figure 3B; Table 2).

248 Phenolic compound

A significant interaction was observed between temperature (18 and 30°C) and pH (7.8 and 7.4) on total phenolic content in *S. capensis* (Factorial ANOVA, $p < 0.01$; Figure 4A). Under both experimental temperatures, total phenolic content increased at pH 7.4 with respect to 7.8.

DISCUSSION

In this investigation, it was observed that the two calcareous macroalgae *L. crispatum* and *S. capensis* suffered similar negative effects subject to the effects of increased temperatures, from 18 (similar to nowadays winter temperatures) to 30°C (comparable to summer end of the 21st Century temperatures), or decreased pH, from current 7.8 to expected future 7.4 (by the year 2100); this was evidenced in terms of photosynthetic performance ($\Delta F/F_m'$), relative carbon content (RCC) and growth rates. However, when increased temperatures (30°C) were combined with acidification (pH 7.4), at least in terms of photosynthesis and growth, both species seemed to be less affected than when both stressors were applied independently. In terms of calcification, acidification at 18°C caused a dramatic decay in RCC in both *L. crispatum* and *S. capensis*. Moreover, although RCC decreased under increased temperatures and acidification, compared to 18°C and pH 7.8, levels remained always higher than at 18°C and 7.4 pH; this for both calcareous macroalgae species. Indeed, increased temperatures appeared to diminish the negative effects induced by acidification, especially in relation to calcification, since at increased temperatures of 30°C and low pH of 7.4 both macroalgae species displayed higher RCC if compared to treatments at 18°C and pH 7.4. Total phenolic content measured in *S. capensis* showed the lowest levels at nowadays temperatures and pH. These increased similarly at 18°C and pH 7.4, and at 30°C and pH 7.8, and with the highest values at 30°C and pH 7.4.

The detrimental effects of increased temperatures on *L. crispatum* and *S. capensis* could be attributed exceeding the tolerance thresholds of both species. Certainly, it has been observed that increased temperatures beyond tolerance ranges can induce an excess in the levels of reactive oxygen species (ROS) and subsequently oxidative stress in other photoautotrophs. For instance, it has been observed that the reaction catalyzed by the

enzyme ribulose-1,5-bisphosphate carboxylase oxygenase (RuBisCO), located in the chloroplast and involved in carbon fixation and photorespiration, can cause greater levels of the ROS H_2O_2 under high temperatures due to increased oxygenase reactions (Kim and Portis 2004). Moreover, it has been observed that increased temperatures induce incomplete H_2O oxidation in the PSII electron donor side, forming H_2O_2 , which is then reduced by manganese to the highly oxidizing $\text{HO}\cdot$ through Fenton reaction (Pospíšil 2016). In this context, Müller et al. (2012) found a systematic increase in the ROS $\cdot\text{O}^{2-}$ in parallel with greater temperatures of up to 18°C in the gametophytes of the brown macroalgae *Alaria esculenta*, *Laminaria digitata*, and *Saccharina latissima*. An extended oxidative stress in macroalgae is capable of causing damage and degradation to photosynthetic pigments; thus, affecting photosynthesis and development (Moenne et al. 2016). The effects of high temperatures have been reported in the rhodolith *Lithophyllum margaritae*, which from 10 to 30°C showed optimum oxygenic photosynthetic threshold at 25°C ; however, when values exceeded these temperatures photosynthesis began to decline (Steller et al. 2007). Reports by Vásquez-Elizondo and Enríquez (2016) demonstrated that from 30°C to 32°C , photosynthesis (F_v/F_m) suffered nearly a 45% decline after 3 d of experiments in rhodolith *Neogoniolithon* sp., articulate *Amphiroa tribulus* and crustace *Lithothamnion* sp. Despite the latter, similar increased-temperature experiments (from optimal) on other tropical calcareous macroalgae, such as *Tricleocarpa cylindrica*, *Padina gymnospora*, and *Lithothamnion corallioides* have not evidenced negative effects in terms of photosynthetic performance (Schermer et al. 2016; Noisette et al. 2013). Therefore, photosynthetic performance under increased temperatures seems to be a species-specific feature in calcifying macroalgae.

In relation to the effects of acidification, our results demonstrated that negative effects of increasing temperatures from 18 to 30°C were softened when decreasing the pH from 7.8 to 7.4 . Our records are similar to those reported by Semesi et al. (2009), which observed that a progressive decrease in pH (from 9.8 to 7.6) were accompanied by an exponential increase in photosynthetic O_2 exchange in the rodolith *Hydrolithon* sp. Recent studies by Scherner et al. (2016) demonstrated the differential effects of low pH on photosynthesis in calcifying macroalgae; while from pH 8.1 to 7.2 F_v/F_m decreased in *Lithophyllum stictaeforme*, *Pneophyllum conicum* and *Porolithon pachydermum*, in

Tricleocarpa cylindrica it increased. Additionally, Kram et al. (2015) observed no change in photosynthesis in the articulated *Jania adhaerens* and in the incrusting *Lithothamnion californicum* from pH 8.1 to 7.6, although growth rates decreased. Even though photosynthesis enhancement has been observed in different macroalgae species subject to acidification, published research suggests that this phenomenon may be more related to CO₂ availability rather than reduction of pH itself. According to Martin and Hall-Spencer (2017), higher concentration of CO₂ in seawater induces macroalgae to rely exclusively of CO₂ diffusion, own-regulating their CO₂-concentrating mechanisms (CCM) to save energy. Certainly, high availability of diffused CO₂ may provide greater substrate for RUBISCO (ribulose-1,5-bisphosphate carboxylase/oxygenase) and carbon fixation, inducing photosynthesis and growth (Martin and Hall-Spencer (2017).

The decalcification of calcifying macroalgae induced by acidification has been already well described (Basso 2012; Feely et al. 2004). Likewise, we observed similar decalcification (about 80%) under lower pH from 7.8 to 7.4 at 18°C in both *L. crispatum* and *S. capensis*. Semesi et al. (2009) showed that a progressive decrease in pH values from a 10 to 7.5 led to nearly 100% decalcification in *Hydrolithon* sp. Similarly, Noisette et al. (2013) observed decrease in calcification rates from pH 8.1 to 7.7 in *Lithothamnion corallioides*. Interestingly, these authors observed that at an increase in environmental temperatures reduced the impact of pH 7.7 in calcification rates. In this context, other studies on coralline algae have shown higher levels of calcification during summer, when average sea surface temperatures are the highest (Steller et al. 2007; Martin and Gattuso 2009). Despite the latter, there are also investigations that account for the combined negative effects of increased temperatures and acidification on calcification rates, as described for the coralline *Porolithon onkodes* (Anthony et al. 2008; Diaz- Pulido et al. 2012). The information suggests that although temperature softened the detrimental effects of acidification (and increased CO₂) in *L. crispatum* and *S. capensis*, this seems to be an intraspecific feature that cannot be applied to all calcifying macroalgae.

Another feature assessed in *S. capensis* was its phenolic content upon exposure to increased temperatures and decreased pH. Total phenolics increased at nowadays temperatures under acidification, but also at increased temperatures and subject to lower pH. Different investigations have revealed that increased temperatures and acidification

are capable of inducing an oxidative stress condition due to de over-production of ROS by different metabolic pathways (e.g. Celis-Plá et al. 2017; Flores- Molina et al. 2016; Pospíšil 2016). It has been well described the role of phenolic compounds as antioxidants to counteract ROS-excess (Moenne et al. 2016); thus, the records may indicate that *S. capensis* induces the production of phenolic compounds to inactivate ROS and avoid oxidative damage.

Short-term assessment of temperature and acidification conditions on physiological and metabolic features of *L. crispatum* and *S. capensis* reinforce the importance of local stressors to promote local baseline shifts. In addition, similar records on other macroalgae species suggest that short-term responses may be similar to those expressed in a long-term and, thus, may provide a good representation of potential tolerance and survival thresholds under upcoming environmental conditions. For instance, Chen et al. (2017) observed that within 60 min, light-saturated net photosynthetic O₂ evolution rate (NPRm) in the green macroalga *Ulva lactuca* decreased steadily when the pH dropped from 8.2 to 7.5 and, even more, to 6.5. After this time, NPRm decrease stopped and levels were maintained without a trend of change. Indeed, similar NPRm levels were observed in the alga exposed to pHs 7.5 and 6.5 at 60 min, 90 min and 8 days of experiments. Similar records on short-term exposure (within 48 h) to future climate change scenarios have recorded in other macroalgae species belonging to different phylogenetic groups (e.g. Celis-Pla et al. 2014; Cruces et al. 2017; Flores- Molina et al. 2016).

In conclusion, we observed that both macroalgae species, the coralline *L. crispatum* and the aragonite *S. capensis*, displayed similar physiological responses to acidification. Moreover, and considering the interaction of higher sea surface temperatures and acidification, the information suggests that both assessed macroalgae species may be able to cope and develop under expected future marine environmental scenarios mediated by Climate Change.

AWKNOWLEDGEMENTS

We thank financial support for Master studies from Becas Chile Magister 2010 (CONICYT, Chile) to Pamela Muñoz. This work was supported by grants from Boticario Foundation; FAPESC—Foundation Support Research and Innovation in the State of Santa Catarina; CNPq—National Council for Scientific and Technological Development; Capes-Higher Education Personnel Improvement Coordination; REBENTOS—Habitat Monitoring Network Coastal Benthic and ProspeMar-Islands— Sustainable Prospecting in Ocean Islands: Biodiversity, Chemistry, Ecology and Biotechnology; Rede Coral Vivo and REDEALGAS; Brazilian Research Network on Global Climate Change, FINEP/Rede CLIMA Grant Number 01.13.0353-00 and FAPESP—Foundation Support Research and Innovation in the State of São Paulo (FAPESP, 2014/00012-1).

REFERENCES

- Amado-Filho G, Maneveldt G, Manso R, Marins-Rosa B, Pacheco M, Guimarães S (2007) Estructura de los mantos de rodolitos de 4 a 55 metros de profundidad en la costa sur del estado de Espírito Santo, Brasil. *Ciencias Marinas* 33:399-410.
- Anthony KR, Kline DI, Diaz-Pulido G, Dove S, Hoegh-Guldberg O (2008) Ocean acidification causes bleaching and productivity loss in coral reef builders. *Proceedings of the National Academy of Sciences* 105:17442-17446.
- Anthony KRN, Maynard JA, Diaz-Pulido G, Mumby PJ, Marshall PA, Cao L, Hoegh-Guldberg O (2011) Ocean acidification and warming will lower coral reef resilience. *Global Change Biology* 17:1798-1808.
- Baker NR (2008) Chlorophyll Fluorescence: A Probe of Photosynthesis In Vivo. *Annual Review of Plant Biology* 59:89-113.
- Basso D (2012) Carbonate production by calcareous red algae and global change. *Geodiversitas* 34:13-33.
- Bischof K, Rautenberger R (2012) Seaweed responses to environmental stress: reactive oxygen and antioxidative strategies. In: *Seaweed Biology*. Springer, UK, pp 109-132.
- Cancellier-Cechinel K (2013) Caracterização físico-química e oceanográfica da Plataforma Continental Interna de Santa Catarina: Subsídio para a gestão da pesca

- 399 da sardinha-verdadeira (*Sardinella brasiliensis*). Master in Environmental Science
400 and Technology, Universidade do Vale do Itajaí, Itajaí, Brazil, 76
- 401 Carmouze JP (1994) O metabolismo dos ecossistemas aquáticos: fundamentos teóricos,
402 métodos de estudo e análises químicas. Edgard Blucher, Sao Paulo, Brazil.
- 403 Celis-Plá PS, Martínez B, Korbee N, Hall-Spencer JM, Figueroa FL (2017)
404 Photoprotective responses in a brown macroalgae *Cystoseira tamariscifolia* to
405 increases in CO₂ and temperature. *Marine Environmental Research* 130:157-165.
- 406 Celis-Pla PSM, Martinez B, Quintano E, Garcia-Sanchez M, Pedersen A, Navarro NP,
407 Copertino MS, Mangaiyarkarasi N, Mariath R, Figueroa FL, Korbee N (2014)
408 Short-term ecophysiological and biochemical responses of *Cystoseira*
409 *tamariscifolia* and *Ellisolandia elongata* to environmental changes. *Aquatic*
410 *Biology* 22:227-243.
- 411 Cruces E, Flores-Molina MR, Díaz MJ, Huovinen P, Gómez I (2017) Phenolics as
412 photoprotective mechanism against combined action of UV radiation and
413 temperature in the red alga *Gracilaria chilensis*? *Journal of Applied Phycology*.
- 414 Chen BB, Zou DH, Zhu MJ (2017) Growth and photosynthetic responses of *Ulva lactuca*
415 (Ulvales, Chlorophyta) germlings to different pH levels. *Mar Biol Res* 13:351-
416 357.
- 417 Dalinghaus C (2016) Análise da estabilidade da forma em planta e perfil nas praias da
418 Barra da Lagoa, Moçambique e Ingleses, Florianópolis - SC: Aplicações em
419 análise de perigos costeiros. Master in Geograpy, Universidade Federal de Santa
420 Catarina, Florianopolis, Brazil, 200
- 421 Diaz- Pulido G, Anthony K, Kline DI, Dove S, Hoegh- Guldberg O (2012) Interactions
422 between ocean acidification and warming on the mortality and dissolution of
423 coralline algae. *Journal of Phycology* 48:32-39.
- 424 Duarte CM, Hendriks IE, Moore TS, Olsen YS, Steckbauer A, Ramajo L, Carstensen J,
425 Trotter JA, McCulloch M (2013) Is ocean acidification an open-ocean syndrome?
426 Understanding anthropogenic impacts on seawater pH. *Estuaries and Coasts*
427 36:221-236.
- 428 Eggert A (2012) Seaweed responses to temperature. In: *Seaweed biology*. Springer,
429 Berlin, pp 47-66.

- 430 Feely RA, Sabine CL, Lee K, Berelson W, Kleypas J, Fabry VJ, Millero FJ (2004)
 431 Impact of anthropogenic CO₂ on the CaCO₃ system in the oceans. Science
 432 305:362-366.
- 433 Ferreira C, Horta PA, Almeida GM, Zitta CS, de M. Oliveira E, Gueye MBYB,
 434 Rodrigues AC (2015) Anatomical and ultrastructural adaptations of seagrass
 435 leaves: an evaluation of the southern Atlantic groups. Protoplasma 252:3-20.
- 436 Figueiredo MAO, Coutinho R, Villas-Boas AB, Tâmega FTS, Mariath R (2012) Deep-
 437 water rhodolith productivity and growth in the southwestern Atlantic. Journal of
 438 Applied Phycology 24:487-493.
- 439 Figueroa F, Barufi JB, Malta E, Conde-Álvarez R, Nitschke U, Arenas F, Mata M,
 440 Connan S, Abreu M, Marquardt R (2014) Short-term effects of increasing CO₂,
 441 nitrate and temperature on three Mediterranean macroalgae: biochemical
 442 composition. Aquatic Biology 22:177-193.
- 443 Findlay HS, Artioli Y, Moreno Navas J, Hennige SJ, Wicks LC, Huvenne VAI,
 444 Woodward EMS, Roberts JM (2013) Tidal downwelling and implications for the
 445 carbon biogeochemistry of cold-water corals in relation to future ocean
 446 acidification and warming. Global Change Biology 19:2708-2719.
- 447 Flores-Molina MR, Rautenberger R, Munoz P, Huovinen P, Gomez I (2016) Stress
 448 tolerance of the endemic Antarctic brown alga *Desmarestia anceps* to UV
 449 radiation and temperature is mediated by high concentrations of phlorotannins.
 450 Photochemistry and Photobiology.
- 451 Flores- Molina MR, Rautenberger R, Munoz P, Huovinen P, Gomez I (2016) Stress
 452 tolerance of the endemic antarctic brown alga *Desmarestia anceps* to UV radiation
 453 and temperature is mediated by high concentrations of phlorotannins.
 454 Photochemistry and Photobiology 92:455-466.
- 455 Glynn PW (1996) Coral reef bleaching: facts, hypotheses and implications. Global
 456 Change Biology 2:495-509.
- 457 Gouvêa LP, Schubert N, Martins CDL, Sissini M, Ramlov F, Rodrigues ERdO, Bastos
 458 EO, Freire VC, Maraschin M, Carlos Simonassi J, Varela DA, Franco D, Cassano
 459 V, Fonseca AL, Barufi J B, Horta PA (2017) Interactive effects of marine

- 460 heatwaves and eutrophication on the ecophysiology of a widespread and
461 ecologically important macroalga. *Limnology and Oceanography* 62:2056-2075.
- 462 Gruber N, Hauri C, Lachkar Z, Loher D, Frölicher TL, Plattner G-K (2012) Rapid
463 progression of ocean acidification in the California Current System. *Science*
464 337:220-223.
- 465 Hall-Spencer JM, Rodolfo-Metalpa R, Martin S, Ransome E, Fine M, Turner SM,
466 Rowley SJ, Tedesco D, Buia M-C (2008) Volcanic carbon dioxide vents show
467 ecosystem effects of ocean acidification. *Nature* 454:96.
- 468 Harley CD, Anderson KM, Demes KW, Jorve JP, Kordas RL, Coyle TA, Graham MH
469 (2012) Effects of climate change on global seaweed communities. *Journal of*
470 *Phycology* 48:1064-1078.
- 471 Harley CDG, Randall Hughes A, Hultgren KM, Miner BG, Sorte CJB, Thornber CS,
472 Rodriguez LF, Tomanek L, Williams SL (2006) The impacts of climate change in
473 coastal marine systems. *Ecology Letters* 9:228-241.
- 474 Hofmann LC, Yildiz G, Hanelt D, Bischof K (2012) Physiological responses of the
475 calcifying rhodophyte, *Corallina officinalis* (L.), to future CO₂ levels. *Marine*
476 *Biology* 159:783-792.
- 477 Hurd CL, Hepburn CD, Currie KI, Raven JA, Hunter KA (2009) Testing the effects of
478 ocean acidification on algal metabolism: considerations for experimental designs.
479 *Journal of Phycology* 45:1236-1251.
- 480 IPCC (2014) The physical science basis. Working group I contribution to the fifth
481 assessment report of the intergovernmental panel on climate change. Cambridge
482 University Press, Cambridge.
- 483 Jury CP, Whitehead RF, Szmant AM (2010) Effects of variations in carbonate chemistry
484 on the calcification rates of *Madracis auretenra* (= *Madracis mirabilis* sensu
485 Wells, 1973): bicarbonate concentrations best predict calcification rates. *Global*
486 *Change Biology* 16:1632-1644.
- 487 Kim K, Portis AR (2004) Oxygen- dependent H₂O₂ production by Rubisco. *Febs Letters*
488 571:124-128.

- 489 Koch M, Bowes G, Ross C, Zhang X-H (2013) Climate change and ocean acidification
490 effects on seagrasses and marine macroalgae. *Global Change Biology* 19:103-
491 132.
- 492 Kram S, Price N, Donham E, Johnson M, Kelly E, Hamilton S, Smith J (2015) Variable
493 responses of temperate calcified and fleshy macroalgae to elevated p CO₂ and
494 warming. *ICES Journal of Marine Science* 73:693-703.
- 495 Latham H (2008) Temperature stress-induced bleaching of the coralline alga *Corallina*
496 *officinalis*: a role for the enzyme bromoperoxidase. *Bioscience Horizons* 1:104-
497 113.
- 498 Martin S, Gattuso JP (2009) Response of Mediterranean coralline algae to ocean
499 acidification and elevated temperature. *Global Change Biology* 15:2089-2100.
- 500 Martin S, Hall-Spencer JM (2017) Effects of Ocean Warming and Acidification on
501 Rhodolith/Maërl Beds. In: Riosmena-Rodríguez R, Nelson W, Aguirre J (eds)
502 Rhodolith/Maërl Beds: A Global Perspective. Springer International Publishing,
503 Cham, pp 55-85.
- 504 Martins-Pereira ML (2004) Estudo da dinâmica das águas do Canal da Barra, Barra da
505 Lagoa – Florianópolis, SC., Bachelor in Geography, Universidade Federal de
506 Santa Catarina, Florianópolis, Brazil, 148
- 507 Melzner F, Thomsen J, Koeve W, Oschlies A, Gutowska MA, Bange HW, Hansen HP,
508 Körtzinger A (2013) Future ocean acidification will be amplified by hypoxia in
509 coastal habitats. *Marine Biology* 160:1875-1888.
- 510 Millero FJ, Lee K, Roche M (1998) Distribution of alkalinity in the surface waters of the
511 major oceans. *Marine Chemistry* 60:111-130.
- 512 Moenne A, González A, Sáez CA (2016) Mechanisms of metal tolerance in marine
513 macroalgae, with emphasis on copper tolerance in Chlorophyta and Rhodophyta.
514 *Aquatic Toxicology* 176:30-37.
- 515 Müller R, Desel C, Steinhoff FS, Wiencke C, Bischof K (2012) UV- radiation and
516 elevated temperatures induce formation of reactive oxygen species in
517 gametophytes of cold- temperate/Arctic kelps (Laminariales, Phaeophyceae).
518 *Phycological Research* 60:27-36.

- 519 Noisette F, Duong G, Six C, Davoult D, Martin S (2013) Effects of elevated pCO₂ on the
 520 metabolism of a temperate rhodolith *Lithothamnion corallioides* grown under
 521 different temperatures. *Journal of Phycology* 49:746-757.
- 522 Orr JC, Fabry VJ, Aumont O, Bopp L, Doney SC, Feely RA, Gnanadesikan A, Gruber N,
 523 Ishida A, Joos F, Key RM, Lindsay K, Maier-Reimer E, Matear R, Monfray P,
 524 Mouchet A, Najjar RG, Plattner G-K, Rodgers KB, Sabine CL, Sarmiento JL,
 525 Schlitzer R, Slater RD, Totterdell IJ, Weirig M-F, Yamanaka Y, Yool A (2005)
 526 Anthropogenic ocean acidification over the twenty-first century and its impact on
 527 calcifying organisms. *Nature* 437:681-686.
- 528 Packard GC, Boardman TJ (1999) The use of percentages and size-specific indices to
 529 normalize physiological data for variation in body size: wasted time, wasted
 530 effort? *Comparative Biochemistry and Physiology Part A: Molecular &*
 531 *Integrative Physiology* 122:37-44.
- 532 Pascelli C, Riul P, Riosmena-Rodríguez R, Scherner F, Nunes M, Hall-Spencer JM, de
 533 Oliveira EC, Horta P (2013) Seasonal and depth-driven changes in rhodolith bed
 534 structure and associated macroalgae off Arvoredo island (southeastern Brazil).
 535 *Aquatic Botany* 111:62-65.
- 536 Pospíšil P (2016) Production of reactive oxygen species by photosystem II as a response
 537 to light and temperature stress. *Frontiers in Plant Science* 7:1950.
- 538 Randhir R, Shetty P, Shetty K (2002) L-DOPA and total phenolic stimulation in dark
 539 germinated fava bean in response to peptide and phytochemical elicitors. *Process*
 540 *Biochemistry* 37:1247-1256.
- 541 Riul P, Lacouth P, Pagliosa PR, Christoffersen ML, Horta PA (2009) Rhodolith beds at
 542 the easternmost extreme of South America: Community structure of an
 543 endangered environment. *Aquatic Botany* 90:315-320.
- 544 Russell BD, Thompson J-AI, Falkenberg LJ, Connell SD (2009) Synergistic effects of
 545 climate change and local stressors: CO₂ and nutrient-driven change in subtidal
 546 rocky habitats. *Global Change Biology* 15:2153-2162.
- 547 Sáez CA, Roncarati F, Moenne A, Moody AJ, Brown MT (2015) Copper-induced intra-
 548 specific oxidative damage and antioxidant responses in strains of the brown alga

- 549 *Ectocarpus siliculosus* with different pollution histories. *Aquatic Toxicology*
550 159:81-89.
- 551 Scherner F, Pereira CM, Duarte G, Horta PA, e Castro CB, Barufi JB, Pereira SMB
552 (2016) Effects of Ocean Acidification and Temperature Increases on the
553 Photosynthesis of Tropical Reef Calcified Macroalgae. *PloS One* 11:e0154844.
- 554 Schreiber U, Neubauer C (1990) O₂-dependent electron flow, membrane energization
555 and the mechanism of non-photochemical quenching of chlorophyll fluorescence.
556 *Photosynthesis Research* 25:279-293.
- 557 Semesi IS, Kangwe J, Björk M (2009) Alterations in seawater pH and CO₂ affect
558 calcification and photosynthesis in the tropical coralline alga, *Hydrolithon*
559 sp.(Rhodophyta). *Estuarine, Coastal and Shelf Science* 84:337-341.
- 560 Smith A, Sutherland J, Kregting L, Farr T, Winter D (2012) Phylomineralogy of the
561 coralline red algae: correlation of skeletal mineralogy with molecular phylogeny.
562 *Phytochemistry* 81:97-108.
- 563 Steller D, Hernández-Ayón J, Riosmena-Rodríguez R, Cabello-Pasini A (2007) Efecto de
564 la temperatura sobre las tasas de fotosíntesis, crecimiento y calcificación del alga
565 coralina de vida libre *Lithophyllum margaritae*. *Ciencias Marinas* 33:441-456.
- 566 Underwood AJ (1997) *Experiments in ecology: their logical design and interpretation*
567 using analysis of variance. Cambridge University Press.
- 568 Vásquez-Elizondo RM, Enríquez S (2016) Coralline algal physiology is more adversely
569 affected by elevated temperature than reduced pH. *Scientific Reports* 6:19030.
- 570 Wernberg T, Russell BD, Moore PJ, Ling SD, Smale DA, Campbell A, Coleman MA,
571 Steinberg PD, Kendrick GA, Connell SD (2011) Impacts of climate change in a
572 global hotspot for temperate marine biodiversity and ocean warming. *Journal of*
573 *Experimental Marine Biology and Ecology* 400:7-16.
- 574 Zavialov PO, Wainer I, Absy JM (1999) Sea surface temperature variability off southern
575 Brazil and Uruguay as revealed from historical data since 1854. *Journal of*
576 *Geophysical Research: Oceans* 104:21021-21032.

577

578

579

580 **Figures**

581 **Figure 1.** Changes in maximum fluorescence ($\Delta F/Fm'$) emitted by (A) *Lithothamnium*
582 *crispatum* and (B) *Sonderophycus capensis* exposed to increased temperatures and
583 decreased pH for up to 48 h. Data correspond to mean \pm SD ($n=9$). Letters represent
584 statistical differences at 95 % confidence interval (SNK tests, $p < 0.05$).

585

586 **Figure 2.** Changes in relative carbon content (RCC), as a proxy of calcification rates,
587 observed in (A) *Lithothamnium crispatum* and (B) *Sonderophycus capensis* exposed to
588 increased temperature and decreased pH for up to 48 h. Data correspond to mean \pm SD
589 ($n=3$). Letters represent statistical differences at 95% confidence interval (SNK tests, $p <$
590 0.05).

591

592 **Figure 3.** Change in growth rate measured in (A) *Lithothamnium crispatum* and (B)
593 *Sonderophycus capensis* exposed to increased temperature and decreased for up to 48 h.
594 Data correspond to mean \pm SD ($n=3$). Letters represent statistical differences at 95%
595 confidence interval (SNK tests, $p < 0.05$).

596

597 **Figure 4.** Change in total phenolic compound in *Sonderophycus capensis* exposed to
598 increased temperature and decreased pH for up 48 h. Data correspond to means \pm SD
599 ($n=6$). Letters represent statistical differences at 95% confidence interval (SNK tests, $p <$
600 0.05).

601

602

603

604

605

606

Table 1 Summary of the chemical characteristics of experimental seawater, including T°, pH, $p\text{CO}_2$, HCO_3^- and CO_3^{2-} concentration, Ω_{calcite} (Ω_{Ca}) and $\Omega_{\text{aragonite}}$ (Ω_{Ar}).

| T(°C) | pH | $p\text{CO}_2$ (ppm) | HCO_3^- ($\mu\text{mol kg SW}^{-1}$) | CO_3^{2-} ($\mu\text{mol kg SW}^{-1}$) | Ω_{Ca} | Ω_{Ar} |
|-------|-------|-------------------------|--|--|----------------------|----------------------|
| 18±2 | 7.8±0 | 432±14 | 1164±51 | 62.563±3.2 | 1.491±0.07 | 0.96±0.04 |
| 18±2 | 7.4±2 | 2035±19 | 1387±13 | 18.816±1.8 | 0.449±0.04 | 0.290±0.02 |
| 30±2 | 7.4±2 | 2504±23 | 1590±15 | 33.393±3 | 0.811±0.07 | 0.543±0.05 |
| 30±2 | 7.8±0 | 455±1.10 | 1148±0.24 | 95.524±0.6 | 2.319±0.01 | 1.553±0.01 |

Note: $n = 3$ (means \pm SDs). Abbreviations: T (°C) = Temperature, $p\text{CO}_2$ = partial pressure of carbon dioxide, HCO_3^- = bicarbonate, CO_3^{2-} = carbonate, Ω_{Ca} = Calcite saturation, Ω_{Ar} = Aragonite Saturation.

Table 2 Summary table of Factorial ANOVA of the effects of temperature and pH treatment on $\Delta F/Fm'$ and calcification percentage in *Lithothamnion crispatum* and *Sonderophycus capensis*.


| | <i>Lithothamnion crispatum</i> | | | | <i>Sonderophycus capensis</i> | | | |
|----------------------------------|--------------------------------|---------|--------|----------|-------------------------------|---------|--------|----------|
| | <i>df</i> | MS | F | <i>p</i> | <i>df</i> | MS | F | <i>p</i> |
| $\Delta F/Fm'$ | | | | | | | | |
| Temperature (T) | 1 | 0.25 | 110.53 | $p<0.01$ | 1 | 0.11 | 25.940 | $p<0.01$ |
| pH | 1 | 0.25 | 108.05 | $p<0.01$ | 1 | 0.60 | 131.84 | $p<0.01$ |
| T*pH | 1 | 0.20 | 88.84 | $p<0.01$ | 1 | 0.15 | 33.76 | $p<0.01$ |
| Error | 32 | 0.002 | | | 32 | 0.004 | | |
| Calcification | | | | | | | | |
| T | 1 | 13.84 | 0.24 | 0,63 | 1 | 0.49 | 0.01 | 0,911 |
| pH | 1 | 4793.10 | 86.23 | $p<0.01$ | 1 | 5159.99 | 141.58 | $p<0.01$ |
| T*pH | 1 | 3301.41 | 59.40 | $p<0.01$ | 1 | 1474.34 | 40.45 | $p<0.01$ |
| Error | 8 | 55.58 | | | 8 | 36.45 | | |

Note:

$n = 8$ for $\Delta F/Fm'$; $n = 3$ for calcification percentage. Abbreviations: T = Temperature, *df* = degrees of freedom.

651

652




653

654 **Figure 1**

655

656

657



658

659

Figure 2

660


661

662

663

664

665



666

667 **Figure 3**


668

669

670

671

672



673

674

Figure 4

675

676